



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 25, 2026 – 04:18 AM EDT

PDB ID : 1ZX9 / pdb_00001zx9
Title : Crystal Structure of Tn501 MerA
Authors : Dong, A.; Ledwidge, R.; Patel, B.; Fiedler, D.; Falkowski, M.; Zelikova, J.;
Summers, A.O.; Pai, E.F.; Miller, S.M.
Deposited on : 2005-06-07
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	467	

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 3587 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Mercuric reductase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	453	3349	2091	598	647	13	0	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	initiating methionine	UNP P00392

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (CCD ID: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).




Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	185	Total 185	O 185	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Mercuric reductase



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	72.54Å 72.54Å 155.39Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.94 – 1.90 29.94 – 1.90	Depositor EDS
% Data completeness (in resolution range)	50.0 (29.94-1.90) 91.1 (29.94-1.90)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	0.05	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.16 (at 1.89Å)	Xtriage
Refinement program	CNS 1.0	Depositor
R, R_{free}	0.215 , 0.234 0.208 , 0.227	Depositor DCC
R_{free} test set	1570 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å ²)	22.5	Xtriage
Anisotropy	0.390	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 28.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3587	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.83% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.37	0/3400	0.91	12/4628 (0.3%)

There are no bond length outliers.

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	165	THR	N-CA-C	-9.75	95.91	109.95
1	A	41	THR	N-CA-C	8.48	120.44	111.03
1	A	140	ASP	N-CA-C	-7.71	104.00	113.41
1	A	359	SER	N-CA-C	-6.92	99.70	110.42
1	A	98	ALA	N-CA-C	6.46	117.98	111.07
1	A	349	THR	N-CA-C	-6.26	100.66	110.36
1	A	30	VAL	N-CA-C	6.05	116.64	108.17
1	A	371	ASP	N-CA-C	-5.88	100.36	109.24
1	A	201	LYS	N-CA-C	-5.64	99.33	108.41
1	A	443	THR	N-CA-C	-5.29	101.49	109.85
1	A	44	ASN	N-CA-C	5.20	119.53	112.04
1	A	202	VAL	N-CA-C	5.07	116.26	108.71

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3349	0	3367	58	0
2	A	53	0	31	1	0
3	A	185	0	0	5	0
All	All	3587	0	3398	59	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:237:THR:HG23	1:A:254:THR:HG23	1.68	0.74
1:A:446:GLU:O	1:A:450:LEU:HD13	1.87	0.73
1:A:17:MET:HE2	1:A:17:MET:HA	1.72	0.71
1:A:53:MET:HE2	1:A:81:ARG:HH22	1.59	0.68
1:A:238:GLN:H	1:A:254:THR:CG2	2.08	0.66
1:A:426:ARG:HA	1:A:426:ARG:HE	1.60	0.66
1:A:225:ALA:HB1	1:A:390:ARG:NH1	2.11	0.66
1:A:253:THR:HG21	1:A:258:LEU:HD12	1.78	0.65
1:A:53:MET:HE1	1:A:170:LEU:HA	1.80	0.62
1:A:17:MET:HA	1:A:17:MET:CE	2.30	0.61
1:A:53:MET:HE2	1:A:81:ARG:NH2	2.14	0.61
1:A:328:ALA:HA	1:A:331:MET:HE3	1.82	0.61
1:A:237:THR:HA	1:A:254:THR:HG21	1.82	0.60
1:A:225:ALA:HB1	1:A:390:ARG:HH12	1.71	0.55
1:A:351:PRO:HD3	1:A:390:ARG:CZ	2.37	0.55
1:A:115:GLY:HA2	1:A:130:ASN:ND2	2.22	0.54
1:A:24:VAL:HG21	1:A:110:ILE:HG13	1.90	0.53
1:A:185:SER:HB2	3:A:1184:HOH:O	2.09	0.52
1:A:377:LEU:HD21	1:A:411:ALA:HB1	1.92	0.51
1:A:446:GLU:HG2	1:A:450:LEU:CD1	2.40	0.51
1:A:382:ARG:HD2	1:A:442:LEU:O	2.11	0.51
1:A:432:GLN:NE2	1:A:432:GLN:H	2.09	0.51
1:A:147:GLY:HA2	1:A:309:ASP:HB2	1.92	0.50
1:A:227:ARG:HH11	1:A:227:ARG:HG2	1.76	0.50
1:A:294:ASP:OD2	1:A:298:ARG:HB2	2.11	0.50
1:A:123:GLN:HE21	1:A:138:MET:HE1	1.77	0.50
1:A:300:SER:O	1:A:302:PRO:HD3	2.12	0.50
1:A:326:ARG:HH11	1:A:326:ARG:HG3	1.77	0.49
1:A:251:LEU:HD12	1:A:251:LEU:N	2.28	0.49
1:A:412:PRO:HD2	1:A:444:MET:HE3	1.95	0.49
1:A:426:ARG:HA	1:A:426:ARG:NE	2.28	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:347:VAL:HB	1:A:353:VAL:HB	1.95	0.48
1:A:341:THR:HB	1:A:426:ARG:HH22	1.78	0.48
1:A:17:MET:CE	1:A:104:LEU:HD21	2.43	0.48
1:A:311:THR:OG1	1:A:313:GLN:HG3	2.13	0.48
1:A:351:PRO:HD3	1:A:390:ARG:NH1	2.28	0.48
1:A:227:ARG:NH2	3:A:1086:HOH:O	2.41	0.48
1:A:91:ARG:NH2	3:A:1091:HOH:O	2.47	0.47
1:A:33:ILE:HD12	1:A:33:ILE:N	2.29	0.47
1:A:446:GLU:HG2	1:A:450:LEU:HD13	1.97	0.46
1:A:42:CYS:HB3	3:A:1095:HOH:O	2.16	0.46
1:A:214:GLU:HB3	1:A:218:ILE:HD12	1.98	0.46
1:A:17:MET:HE3	1:A:104:LEU:HD21	1.98	0.45
1:A:237:THR:HA	1:A:254:THR:CG2	2.45	0.44
1:A:24:VAL:HG21	1:A:110:ILE:CG1	2.47	0.44
2:A:666:FAD:H9	2:A:666:FAD:H1'1	1.85	0.44
1:A:128:ARG:HG3	1:A:128:ARG:HH11	1.84	0.43
1:A:53:MET:CE	1:A:81:ARG:HH22	2.31	0.43
1:A:397:ILE:HD12	1:A:402:HIS:HA	2.00	0.43
1:A:401:SER:O	1:A:402:HIS:HB2	2.19	0.42
1:A:141:ARG:HH11	1:A:141:ARG:HG2	1.84	0.42
1:A:160:GLU:H	1:A:160:GLU:CD	2.27	0.42
1:A:376:THR:OG1	1:A:378:ASP:OD1	2.34	0.42
1:A:262:LYS:HE3	3:A:1176:HOH:O	2.18	0.42
1:A:205:LEU:N	1:A:205:LEU:HD12	2.36	0.41
1:A:53:MET:CE	1:A:81:ARG:NH2	2.84	0.40
1:A:286:ASN:HD21	1:A:290:ALA:HB3	1.87	0.40
1:A:160:GLU:CD	1:A:160:GLU:N	2.79	0.40
1:A:244:HIS:C	1:A:245:MET:HG3	2.47	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	451/467 (97%)	435 (96%)	15 (3%)	1 (0%)	43	36

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	302	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	345/357 (97%)	336 (97%)	9 (3%)	40	35

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	17	MET
1	A	51	LYS
1	A	160	GLU
1	A	254	THR
1	A	302	PRO
1	A	378	ASP
1	A	399	GLU
1	A	432	GLN
1	A	436	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	123	GLN
1	A	130	ASN
1	A	236	HIS
1	A	244	HIS
1	A	255	HIS
1	A	295	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	303	ASN
1	A	315	GLN
1	A	419	GLN
1	A	432	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	FAD	A	666	-	58,58,58	2.16	16 (27%)	85,89,89	1.51	11 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	A	666	-	-	5/34/50/50	0/6/6/6

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	666	FAD	C4X-N5	6.80	1.45	1.30
2	A	666	FAD	C4A-N3A	5.38	1.44	1.34
2	A	666	FAD	C9A-N10	4.89	1.49	1.41
2	A	666	FAD	C2A-N3A	4.42	1.41	1.33
2	A	666	FAD	C9A-C5X	4.31	1.48	1.41
2	A	666	FAD	C5A-C4A	-4.22	1.31	1.39
2	A	666	FAD	C6-C7	3.06	1.43	1.39
2	A	666	FAD	C1'-C2'	-2.98	1.48	1.52
2	A	666	FAD	C8A-N9A	2.84	1.42	1.37
2	A	666	FAD	C10-N1	2.63	1.38	1.33
2	A	666	FAD	C5'-C4'	-2.55	1.48	1.51
2	A	666	FAD	C2'-C3'	-2.45	1.49	1.53
2	A	666	FAD	O4-C4	2.42	1.28	1.23
2	A	666	FAD	C2B-C3B	-2.12	1.47	1.53
2	A	666	FAD	C1'-N10	-2.06	1.42	1.47
2	A	666	FAD	P-O3P	2.02	1.61	1.59

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	666	FAD	C4A-N9A-C8A	-5.16	100.32	105.74
2	A	666	FAD	C5A-C4A-N9A	5.06	111.32	105.81
2	A	666	FAD	N3A-C4A-N9A	-4.63	119.29	127.17
2	A	666	FAD	C1'-N10-C9A	-2.85	115.10	120.63
2	A	666	FAD	C5X-C9A-N10	-2.81	115.42	117.97
2	A	666	FAD	N3A-C2A-N1A	-2.68	124.53	128.58
2	A	666	FAD	O3P-PA-O1A	-2.48	103.24	110.70
2	A	666	FAD	O2B-C2B-C3B	2.41	119.55	111.82
2	A	666	FAD	O3'-C3'-C4'	-2.25	103.81	108.93
2	A	666	FAD	C9-C9A-N10	2.18	124.78	121.85
2	A	666	FAD	O5B-C5B-C4B	-2.17	101.59	108.99

There are no chirality outliers.

All (5) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	666	FAD	C5B-O5B-PA-O1A
2	A	666	FAD	P-O3P-PA-O1A
2	A	666	FAD	C5B-O5B-PA-O3P
2	A	666	FAD	P-O3P-PA-O2A
2	A	666	FAD	O4B-C4B-C5B-O5B

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	666	FAD	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	453/467 (97%)	0.36	26 (5%)	29 31	16, 25, 43, 51	0

All (26) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	104	LEU	3.5
1	A	258	LEU	3.1
1	A	106	GLY	3.1
1	A	334	GLY	3.0
1	A	274	ARG	3.0
1	A	279	ASP	3.0
1	A	5	VAL	2.9
1	A	213	ARG	2.9
1	A	365	HIS	2.8
1	A	301	ASN	2.8
1	A	302	PRO	2.8
1	A	275	SER	2.5
1	A	456	ASN	2.5
1	A	287	ALA	2.5
1	A	24	VAL	2.4
1	A	29	GLN	2.4
1	A	107	ASN	2.4
1	A	97	HIS	2.4
1	A	245	MET	2.3
1	A	333	GLY	2.3
1	A	112	VAL	2.2
1	A	295	GLN	2.2
1	A	4	PRO	2.2
1	A	390	ARG	2.1
1	A	337	ALA	2.1
1	A	368	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	FAD	A	666	53/53	0.95	0.07	17,20,23,25	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.