



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 26, 2026 – 10:50 AM EDT

PDB ID : 7ZY5 / pdb_00007zy5
Title : Crystal structure of compound 2 bound to CK2alpha
Authors : Brear, P.; Fusco, C.; Atkinson, E.; Rossmann, M.; Francis, N.; Iegre, J.; Hyvonen, M.; Spring, D.
Deposited on : 2022-05-23
Resolution : 1.82 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.82 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1112 (1.82-1.82)
Clashscore	190562	1148 (1.82-1.82)
Ramachandran outliers	187476	1140 (1.82-1.82)
Sidechain outliers	187428	1140 (1.82-1.82)
RSRZ outliers	180081	1112 (1.82-1.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	352	25% 83% 8% 7%
1	B	352	4% 84% 7% 7%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 5927 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Casein kinase II subunit alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	326	Total	C	N	O	S	0	2	0
			2767	1772	486	498	11			
1	B	326	Total	C	N	O	S	0	3	0
			2773	1775	487	500	11			

There are 50 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-22	GLY	-	expression tag	UNP P68400
A	-21	SER	-	expression tag	UNP P68400
A	-20	MET	-	expression tag	UNP P68400
A	-19	ASP	-	expression tag	UNP P68400
A	-18	ILE	-	expression tag	UNP P68400
A	-17	GLU	-	expression tag	UNP P68400
A	-16	PHE	-	expression tag	UNP P68400
A	-15	ASP	-	expression tag	UNP P68400
A	-14	ASP	-	expression tag	UNP P68400
A	-13	ASP	-	expression tag	UNP P68400
A	-12	ALA	-	expression tag	UNP P68400
A	-11	ASP	-	expression tag	UNP P68400
A	-10	ASP	-	expression tag	UNP P68400
A	-9	ASP	-	expression tag	UNP P68400
A	-8	GLY	-	expression tag	UNP P68400
A	-7	SER	-	expression tag	UNP P68400
A	-6	GLY	-	expression tag	UNP P68400
A	-5	SER	-	expression tag	UNP P68400
A	-4	GLY	-	expression tag	UNP P68400
A	-3	SER	-	expression tag	UNP P68400
A	-2	GLY	-	expression tag	UNP P68400
A	-1	SER	-	expression tag	UNP P68400
A	0	GLY	-	expression tag	UNP P68400
A	1	SER	-	expression tag	UNP P68400
A	21	SER	ARG	engineered mutation	UNP P68400

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-22	GLY	-	expression tag	UNP P68400
B	-21	SER	-	expression tag	UNP P68400
B	-20	MET	-	expression tag	UNP P68400
B	-19	ASP	-	expression tag	UNP P68400
B	-18	ILE	-	expression tag	UNP P68400
B	-17	GLU	-	expression tag	UNP P68400
B	-16	PHE	-	expression tag	UNP P68400
B	-15	ASP	-	expression tag	UNP P68400
B	-14	ASP	-	expression tag	UNP P68400
B	-13	ASP	-	expression tag	UNP P68400
B	-12	ALA	-	expression tag	UNP P68400
B	-11	ASP	-	expression tag	UNP P68400
B	-10	ASP	-	expression tag	UNP P68400
B	-9	ASP	-	expression tag	UNP P68400
B	-8	GLY	-	expression tag	UNP P68400
B	-7	SER	-	expression tag	UNP P68400
B	-6	GLY	-	expression tag	UNP P68400
B	-5	SER	-	expression tag	UNP P68400
B	-4	GLY	-	expression tag	UNP P68400
B	-3	SER	-	expression tag	UNP P68400
B	-2	GLY	-	expression tag	UNP P68400
B	-1	SER	-	expression tag	UNP P68400
B	0	GLY	-	expression tag	UNP P68400
B	1	SER	-	expression tag	UNP P68400
B	21	SER	ARG	engineered mutation	UNP P68400

- Molecule 2 is 1-NAPHTHOL (CCD ID: 1NP) (formula: C₁₀H₈O) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			11	10	1		
2	B	1	Total	C	O	0	0
			11	10	1		
2	B	1	Total	C	O	0	0
			11	10	1		

- Molecule 3 is ACETATE ION (CCD ID: ACT) (formula: $C_2H_3O_2^-$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	113	Total	O	0	0
			113	113		
4	B	233	Total	O	0	0
			233	233		

4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	64.74Å 68.68Å 332.88Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	26.65 – 1.82 26.65 – 1.82	Depositor EDS
% Data completeness (in resolution range)	94.5 (26.65-1.82) 94.4 (26.65-1.82)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.66 (at 1.82Å)	Xtriage
Refinement program	REFMAC 7.1.016	Depositor
R, R_{free}	0.199 , 0.219 0.214 , 0.229	Depositor DCC
R_{free} test set	3213 reflections (4.78%)	wwPDB-VP
Wilson B-factor (Å ²)	25.7	Xtriage
Anisotropy	0.508	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 47.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5927	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.41% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ACT, INP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.83	1/2842 (0.0%)	1.12	4/3845 (0.1%)
1	B	0.90	2/2848 (0.1%)	1.10	4/3853 (0.1%)
All	All	0.87	3/5690 (0.1%)	1.11	8/7698 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	4
1	B	0	3
All	All	0	7

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	221	MET	SD-CE	-9.95	1.54	1.79
1	A	221	MET	SD-CE	-7.49	1.60	1.79
1	B	150	MET	SD-CE	-5.26	1.66	1.79

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	314	THR	N-CA-C	-5.98	102.81	110.53
1	B	180	GLU	CB-CG-CD	5.95	122.72	112.60
1	A	314	THR	N-CA-C	-5.88	102.95	110.53
1	A	237	ASP	CA-CB-CG	5.77	118.37	112.60
1	B	124	LEU	CA-C-N	5.44	127.83	120.38
1	B	124	LEU	C-N-CA	5.44	127.83	120.38

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	124	LEU	CA-C-N	5.06	127.77	120.38
1	A	124	LEU	C-N-CA	5.06	127.77	120.38

There are no chirality outliers.

All (7) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	18	HIS	Mainchain
1	A	19[A]	ARG	Mainchain
1	A	19[B]	ARG	Mainchain
1	B	18	HIS	Mainchain
1	B	19[A]	ARG	Mainchain
1	B	19[B]	ARG	Mainchain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2767	0	2705	20	0
1	B	2773	0	2709	15	0
2	A	11	0	7	0	0
2	B	22	0	14	2	0
3	B	8	0	6	0	0
4	A	113	0	0	0	0
4	B	233	0	0	2	0
All	All	5927	0	5441	35	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (35) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:50:TYR:HA	1:A:71:LYS:HD2	1.67	0.75
1:A:75:LYS:HZ2	1:A:79:LYS:HG3	1.57	0.69

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:124:LEU:HG	1:A:128:LEU:HD11	1.77	0.66
1:B:17:THR:HG22	4:B:619:HOH:O	1.96	0.65
1:A:27:GLU:HG2	1:A:76:LYS:HE2	1.80	0.62
1:A:15:VAL:O	1:A:19[B]:ARG:HG3	2.00	0.61
1:B:15:VAL:O	1:B:19[B]:ARG:HG3	2.01	0.61
1:B:221:MET:HE2	2:B:401:1NP:O1	2.02	0.59
1:A:21:SER:HA	1:A:24:TRP:CZ2	2.41	0.56
1:A:75:LYS:NZ	1:A:78:ILE:HB	2.21	0.54
1:A:22:GLU:H	1:A:22:GLU:CD	2.16	0.54
1:A:16:ASN:HA	1:A:19[A]:ARG:HD2	1.89	0.54
1:B:19[A]:ARG:HH12	1:B:150:MET:C	2.15	0.54
1:A:75:LYS:NZ	1:A:79:LYS:HG3	2.24	0.53
1:A:75:LYS:HG3	1:A:76:LYS:H	1.75	0.51
1:A:19[A]:ARG:HH12	1:A:150:MET:C	2.19	0.50
1:B:224:SER:HB2	1:B:230:GLU:HG2	1.92	0.50
1:A:224:SER:HB2	1:A:230:GLU:HG2	1.93	0.49
1:B:21:SER:HA	1:B:24:TRP:CZ2	2.48	0.48
1:B:244:ARG:HD3	4:B:628:HOH:O	2.15	0.47
1:A:21:SER:HA	1:A:24:TRP:CE2	2.51	0.45
1:A:75:LYS:O	1:A:76:LYS:HB3	2.15	0.45
1:B:221:MET:CE	2:B:401:1NP:H8	2.46	0.45
1:A:192:VAL:O	1:A:193:ALA:HB3	2.17	0.45
1:B:21:SER:HA	1:B:24:TRP:CE2	2.51	0.45
1:A:75:LYS:HZ1	1:A:78:ILE:HB	1.82	0.43
1:A:76:LYS:HA	1:A:79:LYS:HB2	2.02	0.42
1:B:121:PHE:HZ	1:B:163:MET:HE1	1.84	0.42
1:B:118:ASN:HD22	1:B:164:ILE:H	1.68	0.42
1:B:121:PHE:CZ	1:B:163:MET:HE1	2.55	0.41
1:A:71:LYS:O	1:A:73:VAL:N	2.49	0.41
1:A:157:VAL:HB	1:A:217:SER:HB2	2.02	0.41
1:B:157:VAL:HB	1:B:217:SER:HB2	2.03	0.41
1:B:137:MET:HE1	1:B:218:LEU:HG	2.03	0.41
1:B:303:LYS:HB3	1:B:313:LEU:HG	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	326/352 (93%)	315 (97%)	11 (3%)	0	100	100
1	B	327/352 (93%)	320 (98%)	7 (2%)	0	100	100
All	All	653/704 (93%)	635 (97%)	18 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	302/319 (95%)	292 (97%)	10 (3%)	33	15
1	B	303/319 (95%)	296 (98%)	7 (2%)	44	28
All	All	605/638 (95%)	588 (97%)	17 (3%)	38	21

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	57	ILE
1	A	74	LYS
1	A	75	LYS
1	A	105	VAL
1	A	194	SER
1	A	230	GLU
1	A	237	ASP
1	A	258	ILE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	282	GLU
1	A	327	VAL
1	B	57	ILE
1	B	73	VAL
1	B	103	ASP
1	B	105	VAL
1	B	230	GLU
1	B	258	ILE
1	B	327	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	168	HIS
1	A	186	GLN
1	A	262	ASN
1	B	29	HIS
1	B	118	ASN
1	B	168	HIS
1	B	186	GLN
1	B	262	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	ACT	B	403	-	3,3,3	1.18	0	3,3,3	1.02	0
2	1NP	B	401	-	12,12,12	0.21	0	16,16,16	0.38	0
2	1NP	A	401	-	12,12,12	0.24	0	16,16,16	0.35	0
3	ACT	B	404	-	3,3,3	1.19	0	3,3,3	0.86	0
2	1NP	B	402	-	12,12,12	0.28	0	16,16,16	0.33	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	1NP	B	401	-	-	-	0/2/2/2
2	1NP	B	402	-	-	-	0/2/2/2
2	1NP	A	401	-	-	-	0/2/2/2

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.


There are no ring outliers.

1 monomer is involved in 2 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	401	1NP	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring


in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




Ligand 1NP A 401




Bond lengths




Bond angles



Torsions



Rings



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	326/352 (92%)	1.37	87 (26%) 1 1	22, 47, 80, 97	2 (0%)
1	B	326/352 (92%)	0.06	15 (4%) 37 43	10, 26, 56, 78	3 (0%)
All	All	652/704 (92%)	0.72	102 (15%) 5 5	10, 35, 74, 97	5 (0%)

All (102) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	70	LEU	5.5
1	A	257	TYR	5.1
1	A	73	VAL	4.7
1	B	50	TYR	4.6
1	B	73	VAL	4.5
1	A	261	TYR	4.3
1	A	258	ILE	4.2
1	A	269	PHE	4.1
1	A	327	VAL	4.0
1	A	50	TYR	4.0
1	A	19[A]	ARG	4.0
1	A	286	HIS	3.9
1	A	5	VAL	3.8
1	A	249	LEU	3.7
1	A	255	TYR	3.7
1	A	282	GLU	3.6
1	A	307	TYR	3.5
1	A	263	ILE	3.5
1	A	53	VAL	3.5
1	A	75	LYS	3.4
1	A	254	LEU	3.4
1	A	281	TRP	3.4
1	A	272	ILE	3.4
1	A	292	LEU	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	45	LEU	3.3
1	A	305	LEU	3.2
1	A	251	THR	3.1
1	A	4	PRO	3.1
1	B	125	TYR	3.0
1	A	298	LEU	3.0
1	A	300	PHE	3.0
1	A	17	THR	3.0
1	A	279	LYS	2.9
1	A	302	ASP	2.9
1	A	259	ASP	2.9
1	A	226	ILE	2.9
1	A	265	LEU	2.8
1	B	33	TRP	2.8
1	A	72	PRO	2.8
1	A	33	TRP	2.8
1	A	291	HIS	2.7
1	A	273	LEU	2.7
1	A	107	ARG	2.7
1	A	290	GLN	2.7
1	A	76	LYS	2.7
1	B	121	PHE	2.7
1	A	48	GLY	2.7
1	A	51	SER	2.6
1	A	289	ASN	2.6
1	A	248	VAL	2.6
1	A	2	SER	2.6
1	A	250	GLY	2.6
1	A	206	TYR	2.5
1	A	284	PHE	2.5
1	A	276	HIS	2.5
1	A	46	GLY	2.5
1	A	267	PRO	2.5
1	B	2	SER	2.5
1	A	216	TRP	2.5
1	B	72	PRO	2.5
1	A	306	ARG	2.4
1	A	128	LEU	2.4
1	A	260	LYS	2.4
1	A	3	GLY	2.4
1	B	59	ILE	2.4
1	A	256	ASP	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	271	ASP	2.3
1	B	74	LYS	2.3
1	A	266	ASP	2.3
1	A	9	ALA	2.3
1	A	297	ALA	2.3
1	A	57	ILE	2.3
1	A	71	LYS	2.3
1	A	122	LYS	2.3
1	B	124	LEU	2.3
1	A	109	PRO	2.2
1	A	227	PHE	2.2
1	A	131	TYR	2.2
1	A	324	PHE	2.2
1	B	49	LYS	2.2
1	A	124	LEU	2.2
1	A	294	SER	2.2
1	A	295	PRO	2.2
1	A	246	ALA	2.2
1	A	325	TYR	2.1
1	B	17	THR	2.1
1	A	233	PHE	2.1
1	A	245	ILE	2.1
1	A	271	ASP	2.1
1	B	120	ASP	2.1
1	A	283	ARG	2.1
1	A	62	ASN	2.1
1	A	202	LEU	2.1
1	A	117	ASN	2.1
1	A	262	ASN	2.1
1	B	126	GLN	2.1
1	A	311[A]	SER	2.0
1	A	203	LEU	2.0
1	A	288	GLU	2.0
1	A	74	LYS	2.0
1	A	8	ARG	2.0
1	A	316	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	1NP	A	401	11/11	0.75	0.16	62,64,66,66	0
2	1NP	B	402	11/11	0.79	0.14	39,42,45,50	0
3	ACT	B	404	4/4	0.83	0.13	50,51,51,51	0
2	1NP	B	401	11/11	0.85	0.10	40,44,46,48	0
3	ACT	B	403	4/4	0.93	0.08	34,35,35,36	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around 1NP B 402:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around 1NP B 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.