



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 12:48 PM UTC

PDB ID : 1ZZ2 / pdb_00001zz2
Title : Two Classes of p38alpha MAP Kinase Inhibitors Having a Common Diphenylether Core but Exhibiting Divergent Binding Modes
Authors : Michelotti, E.L.; Moffett, K.K.; Springman, E.B.; Karpusas, M.
Deposited on : 2005-06-13
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	360	17% 67% 22% 6%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	BOG	A	361	X	-	X	-

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 2872 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Mitogen-activated protein kinase 14.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	337	2716	1744	464	496	12	5	0	0

- Molecule 2 is octyl beta-D-glucopyranoside (CCD ID: BOG) (formula: C₁₄H₂₈O₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	A	1	20	14	6	0	0

- Molecule 3 is N-[3-(4-FLUOROPHENOXY)PHENYL]-4-[(2-HYDROXYBENZYL)AMINO]PIPERIDINE-1-SULFONAMIDE (CCD ID: B11) (formula: C₂₄H₂₆FN₃O₄S).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
			Total	C	F	N	O	S		
3	A	1	33	24	1	3	4	1	1	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	103	Total	O	0	0
			103	103		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Mitogen-activated protein kinase 14



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	64.29Å 74.23Å 74.42Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	500.00 – 2.00 52.56 – 2.00	Depositor EDS
% Data completeness (in resolution range)	(Not available) (500.00-2.00) 94.3 (52.56-2.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.57 (at 1.70Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.256 , 0.290 0.247 , 0.281	Depositor DCC
R_{free} test set	3062 reflections (8.07%)	wwPDB-VP
Wilson B-factor (Å ²)	21.2	Xtriage
Anisotropy	0.089	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 31.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.018 for -h,l,k	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	2872	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.69% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: B11, BOG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.43	0/2779	0.90	5/3773 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	12

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	197	TRP	N-CA-C	-7.34	99.58	110.23
1	A	164	LEU	N-CA-C	6.78	120.21	109.50
1	A	186	ARG	N-CA-C	5.71	117.50	111.28
1	A	40	ALA	N-CA-C	-5.32	101.86	110.32
1	A	168	ASP	N-CA-C	5.25	118.64	111.39

There are no chirality outliers.

All (12) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	148	HIS	Sidechain
1	A	169	PHE	Sidechain
1	A	199	HIS	Sidechain
1	A	200	TYR	Sidechain
1	A	24	TYR	Sidechain

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Group
1	A	258	TYR	Sidechain
1	A	30	VAL	Peptide
1	A	307	TYR	Sidechain
1	A	311	TYR	Sidechain
1	A	348	PHE	Sidechain
1	A	35	TYR	Sidechain
1	A	99	PHE	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2716	0	2721	71	0
2	A	20	0	21	9	0
3	A	33	0	25	1	0
4	A	103	0	0	3	0
All	All	2872	0	2767	72	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

All (72) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:236:LEU:HB3	1:A:263:THR:HG23	1.41	1.02
1:A:259:ILE:HG13	2:A:361:BOG:H8'2	1.61	0.81
1:A:148:HIS:HD2	1:A:150:ASP:H	1.33	0.77
1:A:80:HIS:HD2	1:A:82:ASN:H	1.32	0.76
1:A:192:GLU:HB3	2:A:361:BOG:H3'1	1.74	0.69
1:A:80:HIS:CD2	1:A:82:ASN:H	2.10	0.68
1:A:81:GLU:HG3	1:A:136:ARG:HH12	1.61	0.65
1:A:186:ARG:HB2	1:A:194:MET:HE1	1.78	0.64
1:A:242:PRO:HG2	1:A:259:ILE:HD12	1.80	0.63
1:A:259:ILE:HD13	1:A:259:ILE:O	1.98	0.63
1:A:291:LEU:O	2:A:361:BOG:H2'2	1.98	0.63
1:A:149:ARG:HD2	1:A:172:ALA:O	1.99	0.63

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:131:ILE:HG13	1:A:213:MET:HE3	1.79	0.63
1:A:197:TRP:CH2	1:A:255:ALA:HB1	2.35	0.62
1:A:84:ILE:HG22	1:A:166:ILE:O	1.99	0.61
1:A:195:LEU:HD13	1:A:232:LEU:HD21	1.82	0.60
1:A:148:HIS:CD2	1:A:150:ASP:H	2.19	0.59
1:A:259:ILE:HG13	2:A:361:BOG:C8'	2.32	0.59
1:A:10:ARG:H	1:A:10:ARG:HE	1.50	0.59
1:A:282:VAL:O	1:A:286:GLU:HG3	2.03	0.58
1:A:148:HIS:HE1	1:A:167:LEU:O	1.87	0.58
1:A:59:PHE:CD2	1:A:338:LYS:HE2	2.39	0.57
1:A:63:ILE:O	1:A:67:ARG:HG2	2.05	0.57
1:A:290:VAL:O	1:A:296:ARG:HD2	2.05	0.57
1:A:293:SER:OG	2:A:361:BOG:H1	2.04	0.56
1:A:13:LEU:HD23	1:A:39:CYS:SG	2.48	0.53
1:A:236:LEU:HD21	1:A:259:ILE:HD11	1.90	0.53
1:A:81:GLU:CG	1:A:136:ARG:HH12	2.20	0.53
1:A:241:THR:HA	1:A:263:THR:HG21	1.91	0.51
1:A:148:HIS:HD2	1:A:150:ASP:N	2.07	0.50
1:A:10:ARG:HE	1:A:10:ARG:N	2.09	0.50
1:A:141:ILE:HG13	1:A:142:HIS:N	2.25	0.50
1:A:84:ILE:HD11	3:A:362:B11:C29	2.43	0.49
1:A:195:LEU:HD13	1:A:232:LEU:CD2	2.43	0.49
1:A:109:MET:HA	1:A:109:MET:HE2	1.95	0.49
1:A:272:ASN:O	1:A:275:ILE:HG13	2.12	0.48
1:A:259:ILE:HD13	1:A:259:ILE:C	2.38	0.48
1:A:84:ILE:HG13	1:A:106:THR:OG1	2.14	0.48
1:A:234:LEU:HB3	4:A:384:HOH:O	2.14	0.48
1:A:84:ILE:HG23	1:A:169:PHE:HE2	1.79	0.47
1:A:207:TRP:HA	4:A:412:HOH:O	2.14	0.47
1:A:78:MET:HG3	1:A:169:PHE:CZ	2.49	0.47
1:A:256:ARG:O	1:A:259:ILE:HG22	2.15	0.46
1:A:61:SER:HA	1:A:334:ILE:HD11	1.97	0.46
1:A:274:PHE:O	1:A:275:ILE:C	2.58	0.46
1:A:189:ARG:HB3	1:A:194:MET:HE3	1.97	0.45
1:A:123:THR:H	1:A:126:HIS:CD2	2.34	0.45
1:A:290:VAL:O	1:A:296:ARG:CD	2.64	0.45
1:A:330:ARG:HG3	1:A:332:LEU:HD21	1.99	0.45
1:A:330:ARG:CG	1:A:332:LEU:HD21	2.47	0.45
1:A:57:ARG:O	1:A:64:HIS:HD2	2.00	0.45
1:A:270:PHE:CD1	1:A:286:GLU:HG2	2.51	0.45
1:A:148:HIS:O	1:A:149:ARG:HG2	2.18	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:45:LYS:HE2	1:A:45:LYS:HB3	1.89	0.44
1:A:242:PRO:HB2	1:A:247:LEU:HD21	1.99	0.44
1:A:249:LYS:HE2	2:A:361:BOG:O2	2.16	0.44
1:A:199:HIS:O	2:A:361:BOG:H61	2.17	0.44
1:A:242:PRO:HA	4:A:433:HOH:O	2.16	0.44
1:A:195:LEU:HG	1:A:197:TRP:CZ3	2.52	0.44
1:A:125:ASP:HA	1:A:128:GLN:HG3	2.00	0.43
1:A:225:GLY:HA3	1:A:231:GLN:HB2	2.00	0.43
1:A:123:THR:HG22	1:A:125:ASP:H	1.84	0.42
1:A:148:HIS:CE1	1:A:167:LEU:O	2.71	0.42
1:A:84:ILE:CG2	1:A:169:PHE:HE2	2.32	0.42
1:A:109:MET:HB2	1:A:157:ALA:HB1	2.01	0.42
1:A:122:LEU:HD12	1:A:216:LEU:HD22	2.01	0.41
1:A:195:LEU:HG	1:A:197:TRP:CH2	2.55	0.41
1:A:197:TRP:CD1	2:A:361:BOG:H3'2	2.55	0.41
1:A:242:PRO:CG	1:A:259:ILE:HD12	2.48	0.41
1:A:96:LEU:C	1:A:96:LEU:HD13	2.46	0.41
2:A:361:BOG:H7'1	2:A:361:BOG:H4'1	1.81	0.40
1:A:242:PRO:HB2	1:A:247:LEU:CD2	2.52	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	333/360 (92%)	320 (96%)	10 (3%)	3 (1%)	14 9

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	56	SER
1	A	199	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	275	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	298/320 (93%)	273 (92%)	25 (8%)	10 7

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	10	ARG
1	A	13	LEU
1	A	72	LEU
1	A	81	GLU
1	A	84	ILE
1	A	86	LEU
1	A	87	LEU
1	A	94	ARG
1	A	120	GLN
1	A	128	GLN
1	A	130	LEU
1	A	138	LEU
1	A	151	LEU
1	A	185	THR
1	A	194	MET
1	A	195	LEU
1	A	197	TRP
1	A	241	THR
1	A	253	GLU
1	A	259	ILE
1	A	263	THR
1	A	275	ILE
1	A	296	ARG
1	A	333	LEU
1	A	340	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	14	ASN
1	A	64	HIS
1	A	80	HIS
1	A	120	GLN
1	A	126	HIS
1	A	148	HIS
1	A	196	ASN
1	A	264	GLN
1	A	269	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	BOG	A	361	-	20,20,20	3.12	4 (20%)	25,25,25	2.94	11 (44%)
3	B11	A	362	-	35,36,36	7.26	25 (71%)	46,50,50	1.69	10 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BOG	A	361	-	5/5/5/5	3/11/31/31	0/1/1/1
3	B11	A	362	-	-	1/18/30/30	0/4/4/4

All (29) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	362	B11	O18-S16	27.39	1.73	1.43
3	A	362	B11	O19-S16	26.42	1.72	1.43
2	A	361	BOG	O2-C2	-7.95	1.23	1.43
2	A	361	BOG	O3-C3	-7.85	1.23	1.43
2	A	361	BOG	O4-C4	-7.77	1.23	1.43
3	A	362	B11	O7-C6	-6.05	1.24	1.36
3	A	362	B11	F1-C30	5.44	1.49	1.36
3	A	362	B11	C31-C30	4.74	1.46	1.37
3	A	362	B11	C32-C31	4.49	1.46	1.38
3	A	362	B11	C29-C30	4.49	1.45	1.37
3	A	362	B11	C29-C28	4.41	1.45	1.38
3	A	362	B11	C21-C22	4.22	1.45	1.39
3	A	362	B11	C2-C1	4.21	1.46	1.38
3	A	362	B11	S16-N17	4.16	1.73	1.64
3	A	362	B11	C21-C20	4.07	1.46	1.39
3	A	362	B11	C6-C5	4.07	1.45	1.40
3	A	362	B11	C3-C4	4.07	1.45	1.38
3	A	362	B11	C32-C27	4.01	1.46	1.38
3	A	362	B11	C24-C25	3.95	1.45	1.38
3	A	362	B11	C4-C5	3.86	1.45	1.39
3	A	362	B11	C23-C22	3.85	1.45	1.38
3	A	362	B11	C24-C23	3.83	1.45	1.38
3	A	362	B11	C3-C2	3.79	1.46	1.38
3	A	362	B11	C28-C27	3.77	1.45	1.38
3	A	362	B11	C25-C20	3.64	1.45	1.39
3	A	362	B11	C1-C6	3.63	1.45	1.39
3	A	362	B11	O26-C27	-2.46	1.34	1.39
3	A	362	B11	O26-C22	-2.40	1.34	1.39
2	A	361	BOG	O6-C6	-2.29	1.32	1.42

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	361	BOG	O2-C2-C1	4.92	121.80	110.08
2	A	361	BOG	C3-C4-C5	4.86	119.03	110.23
2	A	361	BOG	O4-C4-C5	4.80	121.14	109.32
2	A	361	BOG	O1-C1-C2	4.68	115.39	108.27
2	A	361	BOG	O3-C3-C4	4.42	120.80	110.38
2	A	361	BOG	O3-C3-C2	4.41	120.77	110.38
2	A	361	BOG	C4-C3-C2	4.33	118.43	110.83
2	A	361	BOG	O2-C2-C3	4.32	120.57	110.38
2	A	361	BOG	O4-C4-C3	4.01	119.83	110.38
3	A	362	B11	C4-C5-C6	3.98	122.49	118.16
3	A	362	B11	O19-S16-O18	-3.75	113.11	119.80
3	A	362	B11	C25-C20-N17	-3.71	112.21	120.02
2	A	361	BOG	C1-C2-C3	3.62	117.63	110.01
3	A	362	B11	C25-C20-C21	3.27	123.60	119.66
3	A	362	B11	C8-C5-C6	-3.21	115.40	119.86
2	A	361	BOG	C1-O5-C5	2.63	118.86	113.72
3	A	362	B11	C3-C4-C5	-2.45	117.29	120.88
3	A	362	B11	C28-C29-C30	2.20	120.64	118.38
3	A	362	B11	C32-C27-C28	2.14	123.28	120.16
3	A	362	B11	C14-N13-C12	2.09	114.52	112.12
3	A	362	B11	C8-N9-C10	2.03	117.20	113.84

All (5) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	A	361	BOG	C1
2	A	361	BOG	C3
2	A	361	BOG	C2
2	A	361	BOG	C4
2	A	361	BOG	C5

All (4) torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
2	A	361	BOG	C2-C1-O1-C1'
2	A	361	BOG	O5-C1-O1-C1'
3	A	362	B11	C14-N13-S16-O18
2	A	361	BOG	C4'-C5'-C6'-C7'

There are no ring outliers.

2 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	361	BOG	9	0
3	A	362	B11	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	337/360 (93%)	0.91	61 (18%) 3 3	12, 24, 55, 71	1 (0%)

All (61) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	197	TRP	9.1
1	A	258	TYR	6.6
1	A	184	ALA	6.6
1	A	352	PRO	6.4
1	A	198	MET	6.1
1	A	185	THR	6.1
1	A	251	SER	6.0
1	A	257	ASN	5.9
1	A	255	ALA	5.6
1	A	254	SER	5.3
1	A	259	ILE	5.0
1	A	160	GLU	4.9
1	A	200	TYR	4.7
1	A	275	ILE	4.7
1	A	252	SER	4.3
1	A	256	ARG	4.2
1	A	253	GLU	4.1
1	A	244	ALA	4.0
1	A	199	HIS	3.9
1	A	162	CYS	3.8
1	A	263	THR	3.7
1	A	218	THR	3.6
1	A	226	THR	3.5
1	A	94	ARG	3.4
1	A	34	ALA	3.4
1	A	10	ARG	3.3
1	A	35	TYR	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	17	ILE	3.2
1	A	195	LEU	3.2
1	A	32	SER	3.1
1	A	100	ASN	3.1
1	A	56	SER	3.1
1	A	30	VAL	3.0
1	A	6	PRO	3.0
1	A	241	THR	2.8
1	A	120	GLN	2.8
1	A	33	GLY	2.7
1	A	13	LEU	2.7
1	A	119	CYS	2.7
1	A	167	LEU	2.7
1	A	186	ARG	2.6
1	A	310	GLN	2.6
1	A	110	GLY	2.5
1	A	225	GLY	2.5
1	A	49	ARG	2.5
1	A	117	VAL	2.4
1	A	196	ASN	2.4
1	A	237	ARG	2.4
1	A	220	ARG	2.3
1	A	67	ARG	2.2
1	A	118	LYS	2.2
1	A	123	THR	2.2
1	A	330	ARG	2.2
1	A	116	ILE	2.1
1	A	27	LEU	2.1
1	A	242	PRO	2.1
1	A	240	GLY	2.1
1	A	245	GLU	2.1
1	A	7	THR	2.1
1	A	36	GLY	2.0
1	A	16	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	BOG	A	361	20/20	0.74	0.24	59,61,64,64	0
3	B11	A	362	33/33	0.92	0.09	19,21,22,23	1

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



Electron density around B11 A 362:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.