



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 09:12 PM UTC

PDB ID : 1ZZS / pdb_00001zzs
Title : Bovine eNOS N368D single mutant with L-N(omega)-Nitroarginine-(4R)-Amino-L-Proline Amide Bound
Authors : Li, H.; Flinspach, M.L.; Igarashi, J.; Jamal, J.; Yang, W.; Gomez-Vidal, J.A.; Litzinger, E.A.; Silverman, R.B.; Poulos, T.L.
Deposited on : 2005-06-14
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3428 (1.86-1.86)
Clashscore	190562	3579 (1.86-1.86)
Ramachandran outliers	187476	3553 (1.86-1.86)
Sidechain outliers	187428	3553 (1.86-1.86)
RSRZ outliers	180081	3429 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	416	3% 86% 10% ..
1	B	416	6% 86% 10% ..

2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 7288 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nitric-oxide synthase, endothelial.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	405	Total	As	C	N	O	S	0	0	0
			3219	1	2048	564	590	16			
1	B	405	Total	As	C	N	O	S	0	0	0
			3226	1	2052	567	590	16			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	100	ARG	CYS	see remark 999	UNP P29473
A	368	ASP	ASN	engineered mutation	UNP P29473
A	384	CAS	CYS	modified residue	UNP P29473
B	100	ARG	CYS	see remark 999	UNP P29473
B	368	ASP	ASN	engineered mutation	UNP P29473
B	384	CAS	CYS	modified residue	UNP P29473

- Molecule 2 is ACETATE ION (CCD ID: ACT) (formula: C₂H₃O₂).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 4 2 2	0	0
2	B	1	Total C O 4 2 2	0	0
2	B	1	Total C O 4 2 2	0	0

- Molecule 3 is ZINC ION (CCD ID: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Zn 1 1	0	0

- Molecule 4 is PROTOPORPHYRIN IX CONTAINING FE (CCD ID: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
4	B	1	Total 43	C 34	Fe 1	N 4	O 4	0	0

- Molecule 5 is 5,6,7,8-TETRAHYDROBIOPTERIN (CCD ID: H4B) (formula: $C_9H_{15}N_5O_3$).



- DP9
-
- Chemical structure of DP9, a complex molecule featuring a central pyrrolidine ring. The ring is substituted with a methyl group (CH₃), a carboxylate group (COO⁻), and a side chain. The side chain includes a carboxylate group (COO⁻), a methyl group (CH₃), and a terminal group consisting of a methyl group (CH₃) and a carboxylate group (COO⁻). The structure is shown in a 3D representation with various labels for atoms and groups.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total	C	N	O	0	0
			23	11	8	4		



Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	B	1	Total	C	N	O	0	0
			23	11	8	4		

- Molecule 7 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			6	3	3		
7	B	1	Total	C	O	0	0
			6	3	3		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	331	Total	O	0	0
			331	331		
8	B	321	Total	O	0	0
			321	321		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Nitric-oxide synthase, endothelial



- Molecule 1: Nitric-oxide synthase, endothelial



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	57.87Å 107.06Å 156.79Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.56 – 1.85 46.56 – 1.85	Depositor EDS
% Data completeness (in resolution range)	97.8 (46.56-1.85) 97.8 (46.56-1.85)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.86 (at 1.81Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.175 , 0.211 0.171 , 0.204	Depositor DCC
R_{free} test set	4430 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	20.7	Xtriage
Anisotropy	0.326	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 44.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7288	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.68% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: H4B, CAS, DP9, HEM, ACT, GOL, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.57	0/3300	0.96	13/4495 (0.3%)
1	B	0.54	0/3307	0.97	15/4504 (0.3%)
All	All	0.56	0/6607	0.97	28/8999 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

There are no bond length outliers.

All (28) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	B	360	MET	N-CA-C	-8.65	95.13	109.24
1	B	284	GLY	N-CA-C	-8.12	100.99	112.60
1	A	360	MET	N-CA-C	-7.62	96.82	109.24
1	B	451	VAL	CA-C-N	7.05	124.81	119.66
1	B	451	VAL	C-N-CA	7.05	124.81	119.66
1	B	243	PHE	N-CA-C	-6.65	99.18	109.76
1	A	335	LEU	N-CA-C	6.60	120.56	108.69
1	A	192	TRP	N-CA-C	6.53	119.23	111.33
1	B	245	ILE	N-CA-C	-6.36	98.59	107.80
1	A	452	PRO	CA-C-N	6.35	126.48	119.87
1	A	452	PRO	C-N-CA	6.35	126.48	119.87
1	B	335	LEU	N-CA-C	6.27	119.97	108.69
1	A	226	LEU	N-CA-C	6.19	118.97	110.24
1	A	243	PHE	N-CA-C	-6.17	99.96	109.76

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	359	TYR	N-CA-C	6.12	119.14	110.50
1	A	245	ILE	N-CA-C	-6.06	99.01	107.80
1	B	168	ARG	N-CA-C	-6.03	102.41	110.55
1	A	361	SER	N-CA-C	6.01	119.45	111.75
1	B	192	TRP	N-CA-C	5.91	119.32	111.75
1	B	361	SER	N-CA-C	5.67	119.01	111.75
1	B	296	GLN	N-CA-C	5.46	117.32	108.41
1	A	475	PHE	N-CA-C	-5.33	101.02	109.76
1	A	297	ALA	N-CA-C	-5.23	103.36	110.36
1	B	439	ALA	N-CA-C	5.22	117.05	111.36
1	B	297	ALA	N-CA-C	-5.19	103.41	110.36
1	B	189	ARG	N-CA-C	5.14	119.04	112.87
1	A	168	ARG	N-CA-C	-5.10	103.67	110.55
1	B	370	CYS	N-CA-C	5.09	119.65	113.38

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	137	TYR	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3219	0	3119	28	0
1	B	3226	0	3129	26	0
2	A	4	0	3	0	0
2	B	8	0	6	1	0
3	A	1	0	0	0	0
4	A	43	0	30	0	0
4	B	43	0	30	0	0
5	A	17	0	15	1	0
5	B	17	0	15	1	0
6	A	23	0	22	0	0
6	B	23	0	21	0	0
7	A	6	0	8	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	B	6	0	8	1	0
8	A	331	0	0	2	0
8	B	321	0	0	4	0
All	All	7288	0	6406	55	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (55) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:126:LEU:HD12	1:A:130:ARG:HE	1.37	0.89
1:A:126:LEU:HD11	1:A:156:GLU:HG2	1.55	0.88
1:B:323:GLU:H	1:B:323:GLU:CD	1.95	0.74
1:B:308:PRO:HB2	1:B:311:LEU:HD13	1.74	0.69
1:B:310:GLU:H	1:B:310:GLU:CD	2.02	0.67
1:A:168:ARG:N	1:A:168:ARG:HD2	2.11	0.64
2:B:862:ACT:H1	8:B:1043:HOH:O	2.01	0.61
1:A:126:LEU:CD1	1:A:130:ARG:HE	2.15	0.57
1:A:378:LEU:HB2	8:A:945:HOH:O	2.06	0.56
1:B:370:CYS:SG	1:B:378:LEU:HD13	2.47	0.55
7:B:881:GOL:H2	8:B:1011:HOH:O	2.07	0.55
1:B:72:ARG:HH21	1:B:81:ILE:CD1	2.21	0.53
1:A:370:CYS:SG	1:A:378:LEU:HD13	2.49	0.53
1:A:89:GLN:HG2	8:A:1178:HOH:O	2.08	0.53
1:A:126:LEU:HD23	1:A:159:VAL:HG11	1.90	0.52
1:A:151:ARG:NH2	1:A:168:ARG:HD3	2.25	0.52
1:B:167:LEU:HG	1:B:348:LEU:HD12	1.92	0.52
1:A:126:LEU:HD12	1:A:130:ARG:NE	2.18	0.51
1:B:378:LEU:HB2	8:B:916:HOH:O	2.12	0.50
1:B:72:ARG:HH21	1:B:81:ILE:HD11	1.77	0.50
1:B:249:GLN:HB2	1:B:252:ARG:HD3	1.94	0.49
1:A:449:TRP:HA	5:A:760:H4B:N1	2.26	0.49
1:A:366:THR:HG21	1:A:454:ILE:HG23	1.95	0.48
1:B:323:GLU:CD	1:B:323:GLU:N	2.68	0.48
1:B:69:LYS:HB2	1:B:69:LYS:NZ	2.28	0.48
1:B:308:PRO:HB3	1:B:310:GLU:OE1	2.15	0.47
1:A:151:ARG:CZ	1:A:168:ARG:HD3	2.45	0.47
1:A:409:LEU:O	1:A:413:GLN:HG3	2.16	0.46
1:B:340:ASN:HD22	1:B:340:ASN:H	1.62	0.46
1:B:366:THR:O	1:B:370:CYS:HB2	2.16	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:129:ALA:O	1:A:133:ILE:HG12	2.16	0.46
1:A:275:LEU:O	1:A:279:HIS:HD2	2.00	0.45
1:A:249:GLN:HB2	1:A:252:ARG:CG	2.47	0.45
1:A:340:ASN:H	1:A:340:ASN:HD22	1.64	0.44
1:B:476:ARG:HG2	8:B:1193:HOH:O	2.17	0.44
1:A:130:ARG:CZ	1:A:156:GLU:OE1	2.65	0.44
1:B:449:TRP:HA	5:B:761:H4B:N1	2.32	0.44
1:A:99:ARG:NH2	1:B:88:ALA:O	2.51	0.44
1:B:69:LYS:HB2	1:B:70:PHE:H	1.59	0.44
1:A:168:ARG:N	1:A:168:ARG:CD	2.79	0.43
1:B:472:SER:HA	1:B:473:PRO:C	2.45	0.42
1:A:285:ASN:H	1:A:285:ASN:HD22	1.68	0.42
1:B:340:ASN:HD22	1:B:340:ASN:N	2.17	0.42
1:B:72:ARG:HE	1:B:81:ILE:HD13	1.85	0.42
1:B:231:THR:O	1:B:353:ALA:HA	2.20	0.42
1:A:340:ASN:HD22	1:A:340:ASN:N	2.18	0.41
1:B:266:ASP:HA	1:B:267:PRO:HD3	1.95	0.41
1:A:390:ARG:HE	1:A:390:ARG:HB2	1.75	0.41
1:B:455:SER:HB3	1:B:458:LEU:HD12	2.01	0.41
1:A:231:THR:O	1:A:353:ALA:HA	2.21	0.41
1:A:472:SER:HA	1:A:473:PRO:C	2.45	0.41
1:B:278:GLN:O	1:B:278:GLN:HG2	2.20	0.41
1:B:381:VAL:HG21	1:B:404:ILE:HD11	2.02	0.41
1:A:180:TRP:CE3	1:A:192:TRP:HA	2.56	0.41
1:A:218:LYS:HG2	1:A:311:LEU:HD22	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	400/416 (96%)	391 (98%)	9 (2%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	400/416 (96%)	391 (98%)	8 (2%)	1 (0%)	36	25
All	All	800/832 (96%)	782 (98%)	17 (2%)	1 (0%)	48	35

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	260	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	343/353 (97%)	335 (98%)	8 (2%)	44	30
1	B	344/353 (98%)	339 (98%)	5 (2%)	57	47
All	All	687/706 (97%)	674 (98%)	13 (2%)	50	38

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	126	LEU
1	A	149	GLU
1	A	168	ARG
1	A	259	GLN
1	A	285	ASN
1	A	328	LEU
1	A	340	ASN
1	A	378	LEU
1	B	69	LYS
1	B	260	ASP
1	B	318	GLU
1	B	340	ASN
1	B	378	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (22) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	89	GLN
1	A	153	GLN
1	A	191	GLN
1	A	258	GLN
1	A	259	GLN
1	A	279	HIS
1	A	285	ASN
1	A	340	ASN
1	A	376	ASN
1	A	437	GLN
1	A	468	ASN
1	A	478	GLN
1	B	124	GLN
1	B	146	GLN
1	B	153	GLN
1	B	207	GLN
1	B	222	ASN
1	B	279	HIS
1	B	340	ASN
1	B	405	ASN
1	B	410	HIS
1	B	413	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	CAS	A	384	1	5,8,9	2.95	2 (40%)	1,9,11	0.45	0
1	CAS	B	384	1	5,8,9	2.85	2 (40%)	1,9,11	0.67	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CAS	A	384	1	-	0/0/7/9	-
1	CAS	B	384	1	-	0/0/7/9	-

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	384	CAS	AS-CE2	-4.92	1.83	1.96
1	B	384	CAS	AS-CE2	-4.50	1.84	1.96
1	B	384	CAS	AS-CE1	-4.36	1.84	1.96
1	A	384	CAS	AS-CE1	-4.24	1.84	1.96

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 1 is monoatomic - leaving 11 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	ACT	B	861	-	3,3,3	1.07	0	3,3,3	0.89	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
7	GOL	A	880	-	5,5,5	0.30	0	5,5,5	0.27	0
5	H4B	B	761	-	17,18,18	1.54	3 (17%)	14,26,26	2.08	4 (28%)
4	HEM	B	700	1	50,50,50	1.13	2 (4%)	67,82,82	1.40	8 (11%)
6	DP9	A	799	-	17,23,23	1.04	1 (5%)	20,30,30	1.67	5 (25%)
2	ACT	B	862	-	3,3,3	0.97	0	3,3,3	0.84	0
2	ACT	A	860	-	3,3,3	0.99	0	3,3,3	0.75	0
5	H4B	A	760	-	17,18,18	1.81	4 (23%)	14,26,26	1.94	4 (28%)
4	HEM	A	700	1	50,50,50	1.24	5 (10%)	67,82,82	1.15	5 (7%)
7	GOL	B	881	-	5,5,5	0.23	0	5,5,5	0.20	0
6	DP9	B	800	-	17,23,23	1.06	1 (5%)	20,30,30	1.78	5 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	GOL	A	880	-	-	2/4/4/4	-
5	H4B	B	761	-	-	0/8/17/17	0/2/2/2
6	DP9	A	799	-	-	2/20/32/32	0/1/1/1
4	HEM	B	700	1	-	1/14/54/54	-
5	H4B	A	760	-	-	0/8/17/17	0/2/2/2
4	HEM	A	700	1	-	0/14/54/54	-
7	GOL	B	881	-	-	2/4/4/4	-
6	DP9	B	800	-	-	2/20/32/32	0/1/1/1

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	760	H4B	C4A-N5	4.33	1.48	1.37
5	B	761	H4B	C4A-N5	4.00	1.47	1.37
5	A	760	H4B	C6-N5	3.61	1.54	1.47
5	B	761	H4B	C6-N5	3.27	1.53	1.47
6	B	800	DP9	CG'-N2'	3.27	1.53	1.46
6	A	799	DP9	CG'-N2'	3.13	1.53	1.46
5	A	760	H4B	C7-C6	2.85	1.54	1.52
4	A	700	HEM	FE-ND	2.75	2.03	1.94
4	B	700	HEM	CMA-C3A	2.53	1.56	1.50
4	A	700	HEM	CMA-C3A	2.51	1.55	1.50
4	A	700	HEM	FE-NC	2.32	2.02	1.95

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	700	HEM	C1B-NB	-2.25	1.36	1.40
5	B	761	H4B	C4-N3	2.19	1.43	1.38
4	B	700	HEM	C1D-ND	2.18	1.43	1.38
5	A	760	H4B	C9-C10	2.17	1.57	1.53
4	A	700	HEM	CAB-C3B	-2.16	1.41	1.47

All (31) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	761	H4B	C2-N1-C8A	5.37	122.83	113.36
5	A	760	H4B	C2-N1-C8A	5.07	122.30	113.36
4	B	700	HEM	C3B-C4B-NB	4.46	112.67	109.47
4	B	700	HEM	C4C-C3C-C2C	-4.09	103.27	106.81
6	B	800	DP9	CD-NE-CZ	3.72	130.27	123.46
4	B	700	HEM	CBA-CAA-C2A	-3.56	102.70	112.53
4	B	700	HEM	C4D-ND-C1D	-3.38	101.21	105.21
5	B	761	H4B	O10-C10-C9	-3.34	104.24	109.77
6	B	800	DP9	O'-C'-N1'	-3.33	117.16	123.04
6	A	799	DP9	O'-C'-N1'	-3.27	117.26	123.04
6	A	799	DP9	C'-CA'-N'	-3.16	104.65	111.16
4	A	700	HEM	C3B-C4B-NB	3.14	111.72	109.47
5	A	760	H4B	N3-C2-N1	-3.10	117.64	123.32
6	B	800	DP9	C'-CA'-N'	-3.03	104.92	111.16
5	B	761	H4B	N3-C2-N1	-2.99	117.86	123.32
5	A	760	H4B	O10-C10-C9	-2.76	105.20	109.77
6	A	799	DP9	CD-NE-CZ	2.72	128.44	123.46
6	B	800	DP9	CA'-C'-N1'	2.64	121.26	116.76
4	A	700	HEM	C4B-C3B-C2B	-2.60	104.89	107.28
6	A	799	DP9	CA'-C'-N1'	2.50	121.02	116.76
6	B	800	DP9	CG'-N2'-C	2.46	127.26	123.29
5	B	761	H4B	O9-C9-C6	2.40	115.04	109.28
6	A	799	DP9	CG'-N2'-C	2.33	127.04	123.29
4	A	700	HEM	CMA-C3A-C4A	2.29	128.91	125.42
4	A	700	HEM	C4A-C3A-C2A	-2.28	104.20	106.82
4	B	700	HEM	C3D-C4D-ND	2.21	112.60	110.17
4	A	700	HEM	C3B-C2B-C1B	2.21	108.07	106.41
4	B	700	HEM	CBD-CAD-C3D	-2.17	106.54	112.53
5	A	760	H4B	O9-C9-C6	2.16	114.46	109.28
4	B	700	HEM	C1B-NB-C4B	-2.15	102.66	105.21
4	B	700	HEM	C3C-C2C-C1C	2.09	109.02	107.05

There are no chirality outliers.

All (9) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
7	A	880	GOL	O1-C1-C2-C3
7	B	881	GOL	O1-C1-C2-C3
7	A	880	GOL	O1-C1-C2-O2
7	B	881	GOL	O1-C1-C2-O2
6	A	799	DP9	N-CA-CB-CG
6	B	800	DP9	N-CA-CB-CG
6	B	800	DP9	C-CA-CB-CG
6	A	799	DP9	C-CA-CB-CG
4	B	700	HEM	C4B-C3B-CAB-CBB


There are no ring outliers.

4 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	761	H4B	1	0
2	B	862	ACT	1	0
5	A	760	H4B	1	0
7	B	881	GOL	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	404/416 (97%)	0.04	13 (3%) 50 54	11, 21, 42, 62	0
1	B	404/416 (97%)	0.19	24 (5%) 28 31	11, 24, 44, 62	0
All	All	808/832 (97%)	0.11	37 (4%) 37 40	11, 22, 42, 62	0

All (37) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	119	PRO	6.6
1	B	120	PRO	4.7
1	B	259	GLN	4.7
1	B	121	PRO	4.5
1	A	121	PRO	4.0
1	B	122	ALA	3.4
1	A	238	PRO	3.3
1	B	123	GLU	3.2
1	A	122	ALA	3.1
1	B	311	LEU	3.0
1	A	108	PRO	3.0
1	A	259	GLN	2.8
1	A	239	GLY	2.7
1	B	323	GLU	2.6
1	B	92	GLN	2.6
1	B	131	ASP	2.6
1	B	143	SER	2.5
1	B	138	SER	2.4
1	A	99	ARG	2.4
1	B	260	ASP	2.4
1	B	327	ALA	2.4
1	A	120	PRO	2.4
1	A	157	ALA	2.4
1	B	257	ARG	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	153	GLN	2.3
1	B	326	ALA	2.3
1	B	69	LYS	2.3
1	A	260	ASP	2.2
1	B	310	GLU	2.2
1	B	149	GLU	2.2
1	B	147	ALA	2.2
1	A	261	GLY	2.1
1	A	237	ALA	2.1
1	B	386	ASP	2.1
1	B	156	GLU	2.1
1	B	239	GLY	2.0
1	A	434	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	CAS	B	384	9/10	0.80	0.24	36,39,77,80	0
1	CAS	A	384	9/10	0.91	0.14	20,22,56,64	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	ACT	B	862	4/4	0.70	0.17	52,55,57,58	0
2	ACT	B	861	4/4	0.80	0.18	42,48,49,50	0
7	GOL	B	881	6/6	0.86	0.16	38,44,45,51	0

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
7	GOL	A	880	6/6	0.87	0.15	34,43,45,47	0
2	ACT	A	860	4/4	0.88	0.10	37,39,41,43	0
6	DP9	A	799	23/23	0.90	0.10	20,29,43,44	0
6	DP9	B	800	23/23	0.90	0.12	22,29,39,45	0
5	H4B	A	760	17/17	0.98	0.04	11,13,16,17	0
5	H4B	B	761	17/17	0.98	0.04	9,12,16,17	0
4	HEM	A	700	43/43	0.99	0.05	10,13,21,25	0
4	HEM	B	700	43/43	0.99	0.05	11,14,21,25	0
3	ZN	A	900	1/1	1.00	0.01	18,18,18,18	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around DP9 A 799:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around DP9 B 800:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around HEM A 700:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers ⓘ

There are no such residues in this entry.