



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 8, 2026 – 05:39 PM UTC

PDB ID : 5ZZJ / pdb_00005zzj
Title : Crystal structure of a enzyme from Santalum album
Authors : Han, X.; Liu, W.D.; Chen, C.C.; Guo, R.T.
Deposited on : 2018-06-01
Resolution : 2.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4008 (2.60-2.60)
Clashscore	190562	4347 (2.60-2.60)
Ramachandran outliers	187476	4277 (2.60-2.60)
Sidechain outliers	187428	4277 (2.60-2.60)
RSRZ outliers	180081	4008 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	569	63% 22% 6% 8%
1	B	569	2% 60% 26% 5% 9%
1	C	569	3% 60% 25% 6% 9%
1	D	569	3% 54% 29% 8% 8%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 17019 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Santalene synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	522	Total	C	N	O	S	0	0	0
			4231	2736	707	769	19			
1	B	520	Total	C	N	O	S	0	0	0
			4215	2728	704	764	19			
1	C	519	Total	C	N	O	S	0	0	0
			4211	2726	703	763	19			
1	D	521	Total	C	N	O	S	0	0	0
			4227	2734	706	768	19			


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	45	Total	O	0	0
			45	45		
2	B	45	Total	O	0	0
			45	45		
2	C	35	Total	O	0	0
			35	35		
2	D	10	Total	O	0	0
			10	10		


3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Santalene synthase




• Molecule 1: Santalene synthase







• Molecule 1: Santalene synthase



• Molecule 1: Santalene synthase





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	122.51Å 155.11Å 164.25Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	25.00 – 2.60 25.00 – 2.60	Depositor EDS
% Data completeness (in resolution range)	98.9 (25.00-2.60) 99.2 (25.00-2.60)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.43 (at 2.60Å)	Xtriage
Refinement program	REFMAC 5.8.0222	Depositor
R, R_{free}	0.189 , 0.234 0.217 , 0.250	Depositor DCC
R_{free} test set	4938 reflections (5.10%)	wwPDB-VP
Wilson B-factor (Å ²)	59.5	Xtriage
Anisotropy	0.017	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 34.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	17019	wwPDB-VP
Average B, all atoms (Å ²)	66.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.63% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.17	3/4341 (0.1%)	1.25	13/5879 (0.2%)
1	B	1.14	0/4325	1.28	20/5857 (0.3%)
1	C	1.10	0/4321	1.25	10/5852 (0.2%)
1	D	1.06	1/4337 (0.0%)	1.27	12/5874 (0.2%)
All	All	1.12	4/17324 (0.0%)	1.26	55/23462 (0.2%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	210	LYS	C-N	-7.02	1.24	1.33
1	D	303	PHE	C-N	6.41	1.42	1.33
1	A	409	PRO	C-O	-5.54	1.18	1.24
1	A	383	PRO	C-O	-5.11	1.18	1.24

The worst 5 of 55 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	327	TYR	CA-C-N	-10.47	114.96	122.18
1	C	327	TYR	C-N-CA	-10.47	114.96	122.18
1	B	327	TYR	CA-C-N	-9.40	115.70	122.18
1	B	327	TYR	C-N-CA	-9.40	115.70	122.18
1	A	448	LEU	CA-C-N	8.51	128.45	119.85

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4231	0	4176	114	0
1	B	4215	0	4166	148	0
1	C	4211	0	4161	153	0
1	D	4227	0	4173	197	0
2	A	45	0	0	3	0
2	B	45	0	0	2	0
2	C	35	0	0	2	0
2	D	10	0	0	1	0
All	All	17019	0	16676	605	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 18.

The worst 5 of 605 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:162:CYS:HB3	1:B:195:CYS:SG	1.47	1.50
1:C:141:PHE:O	1:C:142:ILE:HD13	1.30	1.26
1:D:76:PHE:CD2	1:D:104:GLU:HB3	1.71	1.25
1:B:162:CYS:CB	1:B:195:CYS:SG	2.31	1.17
1:C:208:LEU:O	1:C:212:VAL:HG23	1.45	1.16

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	518/569 (91%)	503 (97%)	14 (3%)	1 (0%)	43 66
1	B	516/569 (91%)	503 (98%)	10 (2%)	3 (1%)	21 42
1	C	515/569 (90%)	497 (96%)	17 (3%)	1 (0%)	43 66
1	D	517/569 (91%)	491 (95%)	24 (5%)	2 (0%)	30 51

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	2066/2276 (91%)	1994 (96%)	65 (3%)	7 (0%)	36	58

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	240	ALA
1	B	115	ASP
1	C	164	ASN
1	D	284	ARG
1	D	55	ASN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	459/500 (92%)	393 (86%)	66 (14%)	3	6
1	B	457/500 (91%)	408 (89%)	49 (11%)	6	14
1	C	457/500 (91%)	388 (85%)	69 (15%)	3	5
1	D	459/500 (92%)	382 (83%)	77 (17%)	2	4
All	All	1832/2000 (92%)	1571 (86%)	261 (14%)	3	6

5 of 261 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	330	ILE
1	D	369	VAL
1	D	553	LYS
1	B	391	GLU
1	B	371	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 14 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	61	HIS
1	C	137	GLN
1	D	506	ASN
1	D	98	ASN
1	D	453	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	522/569 (91%)	-0.19	7 (1%) 75 71	35, 55, 91, 134	0
1	B	520/569 (91%)	-0.11	13 (2%) 58 52	30, 57, 95, 134	0
1	C	519/569 (91%)	0.05	15 (2%) 53 48	34, 66, 108, 125	0
1	D	521/569 (91%)	0.27	17 (3%) 49 43	30, 71, 107, 138	0
All	All	2082/2276 (91%)	0.01	52 (2%) 58 52	30, 63, 101, 138	0

The worst 5 of 52 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	446	HIS	7.9
1	C	478	LEU	5.7
1	B	284	ARG	5.3
1	C	110	PHE	5.0
1	B	162	CYS	4.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.