



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 04:15 AM UTC

PDB ID : 5ZZU / pdb_00005zzu
Title : Crystal structure of the C-terminal periplasmic domain of EcEptC from Escherichia coli- complex with Zn
Authors : Zhao, Y.Q.; Cheng, W.; Gu, Y.J.
Deposited on : 2018-06-04
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6658 (2.10-2.10)
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)
RSRZ outliers	180081	6662 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	394	% 86% 10%
1	B	394	2% 82% 9% 9%
1	C	394	% 83% 6% 10%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 9344 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Phosphoethanolamine transferase EptC.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	353	Total	C	N	O	Se	0	0	0
			2882	1827	490	557	8			
1	B	357	Total	C	N	O	Se	0	0	0
			2912	1843	498	564	7			
1	C	353	Total	C	N	O	Se	0	0	0
			2882	1827	490	557	8			

There are 63 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	184	MSE	-	initiating methionine	UNP P0CB39
A	185	GLY	-	expression tag	UNP P0CB39
A	186	SER	-	expression tag	UNP P0CB39
A	187	SER	-	expression tag	UNP P0CB39
A	188	HIS	-	expression tag	UNP P0CB39
A	189	HIS	-	expression tag	UNP P0CB39
A	190	HIS	-	expression tag	UNP P0CB39
A	191	HIS	-	expression tag	UNP P0CB39
A	192	HIS	-	expression tag	UNP P0CB39
A	193	HIS	-	expression tag	UNP P0CB39
A	194	SER	-	expression tag	UNP P0CB39
A	195	SER	-	expression tag	UNP P0CB39
A	196	GLY	-	expression tag	UNP P0CB39
A	197	LEU	-	expression tag	UNP P0CB39
A	198	VAL	-	expression tag	UNP P0CB39
A	199	PRO	-	expression tag	UNP P0CB39
A	200	ARG	-	expression tag	UNP P0CB39
A	201	GLY	-	expression tag	UNP P0CB39
A	202	SER	-	expression tag	UNP P0CB39
A	203	HIS	-	expression tag	UNP P0CB39
A	204	MSE	-	expression tag	UNP P0CB39
B	184	MSE	-	initiating methionine	UNP P0CB39
B	185	GLY	-	expression tag	UNP P0CB39

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	186	SER	-	expression tag	UNP P0CB39
B	187	SER	-	expression tag	UNP P0CB39
B	188	HIS	-	expression tag	UNP P0CB39
B	189	HIS	-	expression tag	UNP P0CB39
B	190	HIS	-	expression tag	UNP P0CB39
B	191	HIS	-	expression tag	UNP P0CB39
B	192	HIS	-	expression tag	UNP P0CB39
B	193	HIS	-	expression tag	UNP P0CB39
B	194	SER	-	expression tag	UNP P0CB39
B	195	SER	-	expression tag	UNP P0CB39
B	196	GLY	-	expression tag	UNP P0CB39
B	197	LEU	-	expression tag	UNP P0CB39
B	198	VAL	-	expression tag	UNP P0CB39
B	199	PRO	-	expression tag	UNP P0CB39
B	200	ARG	-	expression tag	UNP P0CB39
B	201	GLY	-	expression tag	UNP P0CB39
B	202	SER	-	expression tag	UNP P0CB39
B	203	HIS	-	expression tag	UNP P0CB39
B	204	MSE	-	expression tag	UNP P0CB39
C	184	MSE	-	initiating methionine	UNP P0CB39
C	185	GLY	-	expression tag	UNP P0CB39
C	186	SER	-	expression tag	UNP P0CB39
C	187	SER	-	expression tag	UNP P0CB39
C	188	HIS	-	expression tag	UNP P0CB39
C	189	HIS	-	expression tag	UNP P0CB39
C	190	HIS	-	expression tag	UNP P0CB39
C	191	HIS	-	expression tag	UNP P0CB39
C	192	HIS	-	expression tag	UNP P0CB39
C	193	HIS	-	expression tag	UNP P0CB39
C	194	SER	-	expression tag	UNP P0CB39
C	195	SER	-	expression tag	UNP P0CB39
C	196	GLY	-	expression tag	UNP P0CB39
C	197	LEU	-	expression tag	UNP P0CB39
C	198	VAL	-	expression tag	UNP P0CB39
C	199	PRO	-	expression tag	UNP P0CB39
C	200	ARG	-	expression tag	UNP P0CB39
C	201	GLY	-	expression tag	UNP P0CB39
C	202	SER	-	expression tag	UNP P0CB39
C	203	HIS	-	expression tag	UNP P0CB39
C	204	MSE	-	expression tag	UNP P0CB39

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total 1	Zn 1	0	0
2	B	1	Total 1	Zn 1	0	0
2	C	1	Total 1	Zn 1	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	221	Total 221	O 221	0	0
3	B	198	Total 198	O 198	0	0
3	C	246	Total 246	O 246	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Phosphoethanolamine transferase EptC



• Molecule 1: Phosphoethanolamine transferase EptC



• Molecule 1: Phosphoethanolamine transferase EptC



4 Data and refinement statistics

Property	Value	Source
Space group	P 32	Depositor
Cell constants a, b, c, α , β , γ	85.06Å 85.06Å 121.34Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	46.83 – 2.10 46.83 – 2.10	Depositor EDS
% Data completeness (in resolution range)	98.7 (46.83-2.10) 98.7 (46.83-2.10)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.15 (at 2.10Å)	Xtriage
Refinement program	PHENIX (1.11.1_2575: ???)	Depositor
R, R_{free}	0.165 , 0.197 0.165 , 0.198	Depositor DCC
R_{free} test set	2823 reflections (4.92%)	wwPDB-VP
Wilson B-factor (Å ²)	25.7	Xtriage
Anisotropy	0.422	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 33.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.000 for -h,-k,l 0.029 for h,-h-k,-l 0.015 for -k,-h,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	9344	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.86% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.22	0/2953	0.44	0/4011
1	B	0.24	0/2984	0.47	0/4055
1	C	0.25	0/2953	0.46	0/4011
All	All	0.24	0/8890	0.46	0/12077

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2882	0	2757	10	0
1	B	2912	0	2784	18	0
1	C	2882	0	2757	14	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
3	A	221	0	0	1	2
3	B	198	0	0	1	1
3	C	246	0	0	2	0
All	All	9344	0	8298	42	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:356:GLN:HE22	1:B:361:ALA:HB3	1.22	1.02
1:C:251:ARG:NH2	1:C:466:VAL:O	2.24	0.70
1:B:251:ARG:NH1	1:B:430:ASP:OD1	2.33	0.62
1:C:362:ARG:NH2	1:C:363:GLU:OE2	2.33	0.60
1:B:362:ARG:HD3	1:B:364:TYR:OH	2.03	0.59
1:B:268:LEU:HD21	1:B:444:ILE:HG21	1.85	0.58
1:A:314:MSE:HE2	1:A:386:ILE:HD13	1.87	0.55
1:C:249:ARG:NH2	1:C:577:GLN:O	2.37	0.53
1:C:338:THR:HB	3:C:713:HOH:O	2.11	0.51
1:A:296:LEU:HD13	1:A:314:MSE:HE1	1.92	0.50
1:C:505:ARG:HH22	1:C:526:ASP:CG	2.18	0.50
1:A:296:LEU:HD22	1:A:314:MSE:HE1	1.93	0.50
1:B:356:GLN:HE22	1:B:361:ALA:CB	2.09	0.49
1:B:445:LYS:HG3	3:B:768:HOH:O	2.11	0.49
1:B:287:TYR:O	1:B:291:ILE:HG22	2.13	0.48
1:C:251:ARG:NH2	1:C:466:VAL:HB	2.28	0.48
1:C:503:HIS:HE1	3:C:886:HOH:O	1.97	0.47
1:C:287:TYR:O	1:C:291:ILE:HG22	2.14	0.46
1:B:257:TYR:CE2	1:B:259:ARG:HB2	2.50	0.46
1:C:257:TYR:CE2	1:C:259:ARG:HB2	2.50	0.46
1:A:536:ASP:OD2	1:A:539:ARG:HD2	2.16	0.46
1:B:505:ARG:HH22	1:B:526:ASP:CG	2.23	0.46
1:B:500:GLN:HE22	1:B:505:ARG:H	1.64	0.46
1:A:577:GLN:H	1:A:577:GLN:CD	2.24	0.46
1:A:257:TYR:CE2	1:A:259:ARG:HB2	2.51	0.45
1:C:478:ASN:H	1:C:486:MSE:SE	2.51	0.44
1:C:324:THR:HG22	1:C:384:LEU:HB3	2.00	0.43
1:A:404:GLN:NE2	3:A:702:HOH:O	2.43	0.43
1:A:341:THR:O	1:A:345:ARG:HG3	2.18	0.43
1:A:252:MSE:SE	1:A:490:PRO:HB3	2.69	0.43
1:B:252:MSE:SE	1:B:490:PRO:HB3	2.68	0.43
1:C:341:THR:O	1:C:345:ARG:HG3	2.20	0.42
1:B:340:LEU:HD21	1:B:388:HIS:HE1	1.84	0.42
1:C:252:MSE:SE	1:C:490:PRO:HB3	2.70	0.41
1:C:361:ALA:HB3	1:C:398:TYR:HB3	2.01	0.41
1:B:317:MSE:HG3	1:B:384:LEU:HD22	2.02	0.41
1:A:268:LEU:HD21	1:A:444:ILE:HG21	2.02	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:251:ARG:NH2	1:B:466:VAL:HB	2.36	0.41
1:B:277:VAL:HG22	1:B:492:LEU:HD22	2.02	0.41
1:B:478:ASN:H	1:B:486:MSE:SE	2.54	0.40
1:B:359:GLN:HG2	1:B:398:TYR:CD2	2.56	0.40
1:B:365:ASP:O	1:B:368:VAL:HG22	2.22	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:891:HOH:O	3:B:861:HOH:O[1_565]	1.47	0.73
3:A:886:HOH:O	3:A:919:HOH:O[2_554]	1.77	0.43

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	349/394 (89%)	341 (98%)	8 (2%)	0	100	100
1	B	355/394 (90%)	345 (97%)	9 (2%)	1 (0%)	36	36
1	C	349/394 (89%)	341 (98%)	8 (2%)	0	100	100
All	All	1053/1182 (89%)	1027 (98%)	25 (2%)	1 (0%)	48	50

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	357	ARG

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	318/346 (92%)	318 (100%)	0	100	100
1	B	321/346 (93%)	319 (99%)	2 (1%)	78	86
1	C	318/346 (92%)	316 (99%)	2 (1%)	78	86
All	All	957/1038 (92%)	953 (100%)	4 (0%)	84	89

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	451	ASN
1	B	577	GLN
1	C	451	ASN
1	C	577	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	279	ASN
1	A	433	ASN
1	A	436	ASN
1	A	453	ASN
1	B	227	ASN
1	B	279	ASN
1	B	356	GLN
1	B	453	ASN
1	C	227	ASN
1	C	279	ASN
1	C	413	HIS
1	C	453	ASN

5.3.3 RNA

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 3 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	345/394 (87%)	-0.39	3 (0%) 81 83	17, 26, 40, 61	0
1	B	349/394 (88%)	-0.32	6 (1%) 69 71	17, 25, 41, 74	0
1	C	345/394 (87%)	-0.38	2 (0%) 85 87	18, 25, 41, 67	0
All	All	1039/1182 (87%)	-0.36	11 (1%) 78 80	17, 25, 41, 74	0

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	357	ARG	4.9
1	A	361	ALA	4.4
1	C	361	ALA	3.9
1	C	221	ALA	3.8
1	B	334	THR	3.2
1	A	221	ALA	3.0
1	B	337	ASN	2.7
1	B	305	ASP	2.4
1	B	335	ALA	2.3
1	B	336	ARG	2.2
1	A	345	ARG	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	ZN	A	601	1/1	0.95	0.27	64,64,64,64	0
2	ZN	C	601	1/1	0.95	0.27	55,55,55,55	0
2	ZN	B	601	1/1	0.96	0.23	57,57,57,57	0

6.5 Other polymers [i](#)

There are no such residues in this entry.