



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 14, 2026 – 02:17 PM UTC

PDB ID : 7ZZT / pdb_00007zzt
Title : Ligand binding to HDAC2
Authors : Cleasby, A.; Tisi, D.
Deposited on : 2022-05-26
Resolution : 1.56 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.56 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2145 (1.56-1.56)
Clashscore	190562	2189 (1.56-1.56)
Ramachandran outliers	187476	2153 (1.56-1.56)
Sidechain outliers	187428	2150 (1.56-1.56)
RSRZ outliers	180081	2146 (1.56-1.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	498	
1	B	498	
1	C	498	

2 Entry composition

There are 11 unique types of molecules in this entry. The entry contains 10315 atoms, of which 178 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Histone deacetylase 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	369	Total	C	N	O	S	0	9	0
			3015	1929	508	551	27			
1	B	366	Total	C	N	O	S	0	7	0
			2985	1910	500	548	27			
1	C	367	Total	C	N	O	S	0	4	0
			2979	1903	501	549	26			

There are 30 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	493	GLY	-	expression tag	UNP Q92769
A	494	SER	-	expression tag	UNP Q92769
A	495	SER	-	expression tag	UNP Q92769
A	496	GLY	-	expression tag	UNP Q92769
A	497	HIS	-	expression tag	UNP Q92769
A	498	HIS	-	expression tag	UNP Q92769
A	499	HIS	-	expression tag	UNP Q92769
A	500	HIS	-	expression tag	UNP Q92769
A	501	HIS	-	expression tag	UNP Q92769
A	502	HIS	-	expression tag	UNP Q92769
B	493	GLY	-	expression tag	UNP Q92769
B	494	SER	-	expression tag	UNP Q92769
B	495	SER	-	expression tag	UNP Q92769
B	496	GLY	-	expression tag	UNP Q92769
B	497	HIS	-	expression tag	UNP Q92769
B	498	HIS	-	expression tag	UNP Q92769
B	499	HIS	-	expression tag	UNP Q92769
B	500	HIS	-	expression tag	UNP Q92769
B	501	HIS	-	expression tag	UNP Q92769
B	502	HIS	-	expression tag	UNP Q92769
C	493	GLY	-	expression tag	UNP Q92769
C	494	SER	-	expression tag	UNP Q92769
C	495	SER	-	expression tag	UNP Q92769

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	496	GLY	-	expression tag	UNP Q92769
C	497	HIS	-	expression tag	UNP Q92769
C	498	HIS	-	expression tag	UNP Q92769
C	499	HIS	-	expression tag	UNP Q92769
C	500	HIS	-	expression tag	UNP Q92769
C	501	HIS	-	expression tag	UNP Q92769
C	502	HIS	-	expression tag	UNP Q92769

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Zn 1 1	0	0
2	B	1	Total Zn 1 1	0	0
2	C	1	Total Zn 1 1	0	0

- Molecule 3 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Ca 1 1	0	0
3	B	1	Total Ca 1 1	0	0
3	C	1	Total Ca 1 1	0	0

- Molecule 4 is SODIUM ION (CCD ID: NA) (formula: Na).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total Na 1 1	0	0
4	B	1	Total Na 1 1	0	0
4	C	1	Total Na 1 1	0	0

- Molecule 5 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: C₄H₁₀O₃).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	H	O	1	0
			17	4	10	3		
5	A	1	Total	C	H	O	1	0
			17	4	10	3		
5	B	1	Total	C	H	O	1	0
			17	4	10	3		

- Molecule 6 is TRIETHYLENE GLYCOL (CCD ID: PGE) (formula: $C_6H_{14}O_4$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total	C	H	O	1	0
			24	6	14	4		

- Molecule 7 is [(2 {R})-1-oxidanylidene-4-phenyl-1-pyrrolidin-1-yl-butan-2-yl]azanium (CCD ID: KJF) (formula: C₁₄H₂₁N₂O) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
7	A	1	Total	C	H	N	O	0	0
			38	14	21	2	1		
7	B	1	Total	C	H	N	O	0	0
			38	14	21	2	1		
7	C	1	Total	C	H	N	O	0	0
			38	14	21	2	1		

- Molecule 8 is 2-[N-CYCLOHEXYLAMINO]ETHANE SULFONIC ACID (CCD ID: NHE) (formula: C₈H₁₇NO₃S).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
8	B	1	Total	C	H	N	O	S	
			30	8	17	1	3	1	
								0	0

- Molecule 9 is TETRAETHYLENE GLYCOL (CCD ID: PG4) (formula: $C_8H_{18}O_5$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
9	B	1	Total	C	H	O	1	0
			31	8	18	5		
9	B	1	Total	C	H	O	1	0
			31	8	18	5		

- Molecule 10 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
10	B	1	Total 10	C 2	H 6	O 2	1	0
10	B	1	Total 10	C 2	H 6	O 2	1	0
10	C	1	Total 10	C 2	H 6	O 2	1	0


- Molecule 11 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
11	A	333	Total 333	O 333	0	0
11	B	403	Total 403	O 403	0	0
11	C	280	Total 280	O 280	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Histone deacetylase 2



• Molecule 1: Histone deacetylase 2



• Molecule 1: Histone deacetylase 2



GLN	ALA	ILE	PRO	GLU	ASP	ALA	VAL	HIS	GLU	ASP	SER	GLY	ASP	GLU	GLY	GLY	ASP	GLU	GLY	ASP	PRO	ASP	LYS	ARG	LYS	ILE	SER	TYR	ILE	ARG	ASN	ALA	SER	SER	GLY	ASP	LYS	ARG	LYS	THR	ILE	ALA	CYS	ASP	GLU	GLU	PHE	SER	SER	GLU	ASP	SER	GLN	LEU	SER	ASN	PRO	GLY	GLY	SER	SER	GLY	GLY	ARG	ARG	ASN	VAL	HIS	ALA	HIS	HIS	HIS	HIS	LYS	LYS	GLY	ALA
LYS	LYS	ALA	ARG	ILE	GLU	GLU	ASP	LYS	LYS	GLU	THR	GLU	ASP	LYS	LYS	THR	VAL	LYS	GLU	GLU	ASP	ASP	LYS	GLY	LYS	THR	ASP	ASN	ALA	SER	SER	GLY	GLY	LYS	LYS	THR	THR	ASP	LYS	LYS	GLY	THR	LYS	SER	SER	GLU	GLN	LEU	SER	ASN	PRO	GLY	GLY	SER	SER	GLY	GLY	ARG	HIS	HIS	HIS	HIS	HIS	LYS	LYS	GLY	ALA										

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	92.21Å 98.02Å 139.28Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.39 – 1.56 48.39 – 1.56	Depositor EDS
% Data completeness (in resolution range)	98.5 (48.39-1.56) 98.6 (48.39-1.56)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.29 (at 1.55Å)	Xtriage
Refinement program	REFMAC 5.8.0222	Depositor
R, R_{free}	0.148 , 0.173 0.160 , 0.183	Depositor DCC
R_{free} test set	8843 reflections (4.90%)	wwPDB-VP
Wilson B-factor (Å ²)	18.6	Xtriage
Anisotropy	0.496	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 44.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	10315	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.47% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NHE, CA, NA, ZN, EDO, KJF, PG4, PGE, PEG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.77	0/3117	0.92	4/4206 (0.1%)
1	B	0.81	1/3081 (0.0%)	0.93	5/4158 (0.1%)
1	C	0.72	0/3065	0.84	0/4137
All	All	0.77	1/9263 (0.0%)	0.90	9/12501 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3
1	B	0	3
1	C	0	2
All	All	0	8

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	258	GLN	C-O	-5.62	1.21	1.23

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	349	HIS	CA-CB-CG	-6.04	107.76	113.80
1	B	244	GLN	CA-C-N	-5.82	115.27	121.84
1	B	244	GLN	C-N-CA	-5.82	115.27	121.84
1	B	349	HIS	CA-CB-CG	-5.77	108.03	113.80
1	A	320	GLU	CB-CG-CD	5.75	122.38	112.60

There are no chirality outliers.

5 of 8 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	39	ARG	Sidechain
1	A	60	ARG	Sidechain
1	A	98	ARG	Sidechain
1	B	39	ARG	Sidechain
1	B	82	ARG	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3015	0	2946	23	0
1	B	2985	0	2906	17	0
1	C	2979	0	2890	12	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
4	C	1	0	0	0	0
5	A	14	20	20	0	0
5	B	7	10	10	0	0
6	A	10	14	14	0	0
7	A	17	21	0	0	0
7	B	17	21	0	0	0
7	C	17	21	0	0	0
8	B	13	17	16	0	0
9	B	26	36	36	0	0
10	B	8	12	12	0	0
10	C	4	6	6	0	0
11	A	333	0	0	2	0
11	B	403	0	0	2	0
11	C	280	0	0	4	0
All	All	10137	178	8856	52	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 52 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:233[B]:MET:HE1	1:A:237:ILE:HG21	1.51	0.92
1:A:233[B]:MET:HE2	1:A:279:PHE:CZ	2.06	0.90
1:B:245:ILE:HG21	1:B:364[A]:MET:HE3	1.63	0.80
1:C:199:MET:HE1	1:C:378:LEU:HD11	1.73	0.70
1:B:245:ILE:HG22	1:B:368[B]:LYS:HD2	1.74	0.69

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	376/498 (76%)	370 (98%)	6 (2%)	0	100	100
1	B	371/498 (74%)	367 (99%)	4 (1%)	0	100	100
1	C	369/498 (74%)	365 (99%)	4 (1%)	0	100	100
All	All	1116/1494 (75%)	1102 (99%)	14 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	326/425 (77%)	324 (99%)	2 (1%)	78	64
1	B	323/425 (76%)	320 (99%)	3 (1%)	70	51
1	C	321/425 (76%)	319 (99%)	2 (1%)	78	64
All	All	970/1275 (76%)	963 (99%)	7 (1%)	78	59

5 of 7 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	341	TYR
1	B	378	LEU
1	C	366	LYS
1	C	326	ASP
1	B	151	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	95	GLN
1	B	133	GLN
1	C	62	HIS
1	C	244	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 22 ligands modelled in this entry, 9 are monoatomic - leaving 13 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
9	PG4	B	606	-	12,12,12	0.46	0	11,11,11	0.28	0
9	PG4	B	605	-	12,12,12	0.36	0	11,11,11	0.28	0
10	EDO	C	604	-	3,3,3	0.40	0	2,2,2	0.25	0
5	PEG	A	605	-	6,6,6	0.36	0	5,5,5	0.42	0
7	KJF	B	610	2	18,18,18	0.45	0	23,23,23	0.49	0
5	PEG	A	604	-	6,6,6	0.47	0	5,5,5	0.30	0
7	KJF	C	605	2	18,18,18	0.37	0	23,23,23	0.43	0
7	KJF	A	607	2	18,18,18	0.23	0	23,23,23	0.58	0
8	NHE	B	604	-	13,13,13	0.56	0	16,17,17	0.63	0
5	PEG	B	608	-	6,6,6	0.52	0	5,5,5	0.30	0
10	EDO	B	607	-	3,3,3	0.45	0	2,2,2	0.43	0
10	EDO	B	609	-	3,3,3	0.33	0	2,2,2	0.48	0
6	PGE	A	606	-	9,9,9	0.38	0	8,8,8	0.39	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
9	PG4	B	606	-	-	1/10/10/10	-
9	PG4	B	605	-	-	4/10/10/10	-
10	EDO	C	604	-	-	1/1/1/1	-
5	PEG	A	605	-	-	3/4/4/4	-
7	KJF	B	610	2	-	0/13/20/20	0/2/2/2
5	PEG	A	604	-	-	1/4/4/4	-
7	KJF	C	605	2	-	0/13/20/20	0/2/2/2
7	KJF	A	607	2	-	0/13/20/20	0/2/2/2
8	NHE	B	604	-	-	0/7/15/15	0/1/1/1
5	PEG	B	608	-	-	3/4/4/4	-
10	EDO	B	607	-	-	0/1/1/1	-
10	EDO	B	609	-	-	0/1/1/1	-
6	PGE	A	606	-	-	4/7/7/7	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 17 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
6	A	606	PGE	O2-C3-C4-O3
5	A	605	PEG	O1-C1-C2-O2
9	B	605	PG4	O3-C5-C6-O4
9	B	605	PG4	O1-C1-C2-O2
6	A	606	PGE	C4-C3-O2-C2

There are no ring outliers.


No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand KJF B 610



Ligand KJF C 605





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	369/498 (74%)	-0.11	2 (0%) 87 90	10, 21, 39, 55	9 (2%)
1	B	366/498 (73%)	-0.17	1 (0%) 90 93	9, 19, 38, 54	7 (1%)
1	C	367/498 (73%)	0.17	7 (1%) 66 75	14, 26, 44, 72	4 (1%)
All	All	1102/1494 (73%)	-0.04	10 (0%) 81 86	9, 22, 42, 72	20 (1%)

The worst 5 of 10 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	379	PRO	3.4
1	A	11	GLY	3.2
1	C	341	TYR	2.5
1	C	54	ARG	2.4
1	C	344	PRO	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
10	EDO	B	607	4/4	0.77	0.15	36,54,55,58	1
9	PG4	B	605	13/13	0.84	0.16	30,40,61,73	1
5	PEG	A	604	7/7	0.85	0.12	36,46,52,53	1
5	PEG	A	605	7/7	0.85	0.14	36,58,61,65	1
5	PEG	B	608	7/7	0.86	0.12	36,43,56,57	1
6	PGE	A	606	10/10	0.87	0.13	36,47,56,64	1
9	PG4	B	606	13/13	0.91	0.11	30,39,42,45	1
10	EDO	B	609	4/4	0.91	0.10	35,40,46,52	1
10	EDO	C	604	4/4	0.91	0.12	36,46,55,58	1
8	NHE	B	604	13/13	0.92	0.11	20,23,28,33	30
7	KJF	B	610	17/17	0.97	0.05	13,14,16,16	0
7	KJF	C	605	17/17	0.97	0.05	13,15,18,18	38
4	NA	A	603	1/1	0.98	0.04	21,21,21,21	0
4	NA	B	603	1/1	0.98	0.04	20,20,20,20	0
4	NA	C	603	1/1	0.98	0.07	20,20,20,20	0
7	KJF	A	607	17/17	0.99	0.03	13,15,17,18	0
3	CA	B	602	1/1	1.00	0.10	21,21,21,21	0
3	CA	C	602	1/1	1.00	0.13	26,26,26,26	0
2	ZN	A	601	1/1	1.00	0.01	14,14,14,14	0
2	ZN	B	601	1/1	1.00	0.01	13,13,13,13	0
2	ZN	C	601	1/1	1.00	0.02	18,18,18,18	0
3	CA	A	602	1/1	1.00	0.12	21,21,21,21	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around KJF B 610:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around KJF C 605:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around KJF A 607:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.